Magnesium Glycinate

Improved

NPN 80029196

DESCRIPTION

Magnesium Glycinate tablets supply 100 mg of elemental magnesium from a magnesium glycinate complex.

FUNCTIONS

Magnesium is a mineral with a fundamentally important physiological function in the body. However, typical diets in Canada and other industrialized countries often provide less than adequate amounts of magnesium. Clinical evidence indicates that magnesium glycinate has a significantly higher bioavailability than other forms of magnesium, and is less likely to cause a laxative effect. Supplementation with a bioavailable magnesium glycinate complex can help bridge the gap between dietary intake and optimal requirement.

Magnesium plays an essential role in a wide range of fundamental cellular reactions. More than 300 enzymes require magnesium as a cofactor. Complexed with adenosine triphosphate (ATP), the main carrier of metabolic energy in the body, magnesium is essential for all biosynthetic processes: glycolysis, formation of cyclic adenosine monophosphate (cAMP), energy-dependent membrane transport, and muscle function. Two-thirds of the body's magnesium content is located in the skeleton, where it helps to maintain bone health. Recent scientific studies show that magnesium supplementation in animals not only promotes bone formation while increasing its dynamic strength, but also prevents bone resorption.

INDICATIONS

- Muscle function
- Bone health

FORMULA (#202559-120HYC-C)

Each Tablet Contains:

Non-medicinal ingredients: Hyprolose, ascorbyl palmitate, croscarmellose sodium, silica, carnauba wax, tablet coating (polyvinyl alcohol, polyethylene glycol, polysorbate 80, talc).

SUGGESTED USAGE

Adults: Take 1 tablet 1-2 times daily or as directed by your healthcare practitioner.

STORAGE

Store in a cool, dry place. Keep out of reach of children.

Magnesium Glycinate

Improved

NPN 80029196

REFERENCES

Launius BK, Brown PA, Cush EM, Mancini MC. Crit Care Nurs Q. 2004 Jan-Mar;27(1):96-100. Review. Chang JJ, Mack WJ, Saver JL, Sanossian N. Front Neurol. 2014 Apr 15;5:52. doi: 10.3389/fneur.2014.00052. Price CT, Langford JR, Liporace FA. Open Orthop J. 2012;6:143-9. doi: 10.2174/1874325001206010143. Rude RK, Singer FR, Gruber HE. J Am Coll Nutr. 2009 Apr;28(2):131-41. Review. Vormann J. Magnesium: nutrition and metabolism. Mol Aspects Med. 2003 Feb-Jun;24(1-3):27-37. Review. Sun-Edelstein C MA. Expert Rev Neuother. 2009;3(Mar; 9):369-379. Rylander R, Bullarbo M. [304-POS]: Pregnancy Hypertens. 2015 Jan;5(1):150. doi: 10.1016/j.preghy.2014.10.310. João Matias P, et al. Blood Purif. 2014;38(3-4):244-52. doi: 10.1159/000366124. Rodríguez-Moran M, Guerrero-Romero F. Arch Med Res. 2014 Jul;45(5):388-93. doi: 10.1016/j.arcmed.2014.05.003. Blaine J, Chonchol M, Levi M. Clin J Am Soc Nephrol. 2014 Oct 6. pii: CJN.09750913

For more information on Magnesium Glycinate visit douglaslabs.ca

Douglas Laboratories 490 Elgin Mills Road E. Richmond Hill, ON L4C 0L8 866-856-9954 douglaslabs.ca



© 2019 Douglas Laboratories. All Rights Reserved

2